## POISON

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Regiment 550

### FUNGICIDE

#### ACTIVE CONSTITUENT: 550 g/L PROPICONAZOLE SOLVENT: 371 g/L LIQUID HYDROCARBON

## GROUP **3** FUNGICIDE

For the control of certain fungal diseases of Bananas, Oats, Peanuts, Perennial Ryegrass, Pineapples, Stone Fruit, Sugar Cane, Wheat and other crops in certain states and for the control of Dollar spot in Bent and Queensland Blue Couch, and Spring Dead Spot in Couch as specified in the DIRECTIONS FOR USE.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT



Indigo Specialty Products Pty Ltd ABN 15 631 459 660 6/163-173 McEvoy Street, Alexandria NSW 2015 Ph: 0402 725 887 Distributed by





#### (Graferee Regiment 550 FUNGICIDE

#### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. D0 NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. D0 NOT dispose of undiluted chemical on-site. For pineapples dispose of spent dip in approved landfill or bury under at least 500mm of soil in a non-crop, non- pasture area away from water sources or homes. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the nose and throat, and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and a disposable mist face mask to cover the mouth and nose. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and a disposable mist face mask to cover the mouth and nose. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles. If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

#### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed D0 NOT induce vomiting.

#### SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce REGIMENT 550 Fungicide is available from Indigo Specialty Products Pty Ltd on request.

#### Combustible liquid.

In case of fire: Suitable extinguishing media are carbon dioxide, dry chemical, foam, water fog.

#### May cause an allergic skin reaction.

If skin irritation or rash occurs: Get medical advice.

Do not eat, drink or smoke when using this product.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

#### **CONDITIONS OF SALE**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use.

No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



IN A TRANSPORT



Batch No.

Date of Manufacture

APVMA Approval No.: 68606/121766

FLAT SIZE: 332 MM WIDE X 448 MM DEEP

#### FOLDS TO: 83 MM WIDE X 112 MM DEEP

## **Regiment 550**

#### **FUNGICIDE**

#### DIRECTIONS FOR USE: RESTRAINTS

DO NOT apply to turf under heat or moisture stress.

DO NOT apply to more than one application per year on the following couch varieties: C29, Greenless Park, National Park, Tifway, Sportsway and Wagga City.

Crop	Disease Controlled	State	Rate Per Hectare	High volume	WHP	Critical Comments
Apricots	Prune Rust ( <i>Tranzschelia</i> discolour)	SA only	-	Dilute Spraying: 15 mL/ 100L Concentrate Spraying: Refer to the Mixing/ Application Section	1 day	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. <b>Curative Control</b> : Apply when the disease first occurs. Further applications should be made when the disease occurs on new growth. <b>DO NOT</b> make more than 5 applications to any individual tree during the season. <b>Protective Treatment:</b> Spray mancozeb or zineb mixed with REGIMENT 550 at the full recommended rates of application. <b>This use is subject to a DMI anti-resistance strategy</b> .
Bananas (including bananas inter- planted with avocados)	Leaf Spot (Mycosphaerella musicola), Leaf Speckle (Mycosphaerella musae), Cordana Leaf Spot (Cordana johnstonii)	Sth Qld, NSW only	Ground: 90 to 185 mL + 3 to 5L of water miscible oil in a convenient volume of water Aerial: 185 mL + 3 to 5L of a water miscible oil in a minimum of 30L of water Aerial Application without water: 185 mL + 8 to 10L of spraying oil. (This does not require further dilution with water)	-	1 day	This is subject to a DMI anti-resistance strategy. Ground Application: Apply by misting machine or airblast sprayer. Use rates towards the higher end of the range where weather conditions favour diseases or where equipment or terrain does not permit thorough spray coverage of all foliage. Sth Old, NSW: Ground and Aerial application: Commence spraying at the start of the summer rainy season. Apply a maximum of 5 Group 3 sprays per season. Apply a maximum of 2 consecutive sprays of REGIMENT 550 at 21 to 28 day intervals before applying further treatments of a protectant fungicide from a different activity group.
	Leaf Spot (Mycosphaerella musicola), Leaf Speckle (Mycosphaerella musae), Cordana Leaf Spot (Cordana johnstonii) Black Sigatoka (Mycosphaerella fijiensis var difformis)	Nth Qld, WA, NT only Qld, WA, NT only	Ground Application: 185 mL + 3 to 5L of water miscible oil in a convenient volume of water Aerial Application: 185 mL + 3 to 5L of water miscible oil, in a minimum of 30L of water Aerial Application without water: 185 mL + 8 to 10L of spraying oil. (This use does not require further dilution with water)	-	1 day	Nth Qld, WA, NT: Ground and aerial applications: Commence spraying at the start of the wet season. Apply a regular schedule of protectant sprays. When the disease potential is high, apply a maximum of 2 consecutive sprays of REGIMENT 550 at 14 to 21 day intervals before applying further treatments of a protectant fungicide from a different activity group. Apply a maximum of 6 Group 3 sprays per season. Continue with treatments using an alternative protectant fungicide for the remainder of the season. Use the lower rate of oil in Nth Qld. <b>DO NOT</b> apply during July, August, September and October.
Barley	Powdery Mildew ( <i>Blumeria graminis</i> ) Barley Scald ( <i>Rhynchosporium secalis</i> )	All states	70 to 230 mL 230 mL	-	Harvest 4 weeks Grazing 7 days	Spray at the first sign of the disease during the tillering stage (Z21-Z29). A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Powdery Mildew: Higher rates provide longer protection.
	Spot Form Net Blotch (Pyrenophora teres f. maculate)					Apply after flag leaf is around 70% emerged (Z38) and before infection averages 10% on the flag-2 leaf. Ensure thorough coverage of stems and leaves.
Boronia	Rust (Puccinia boroniae)	Tas, WA only	230 to 460 mL	-	-	Apply 2 to 5 applications at 10 to 14 days intervals during the main disease period. Use the lower rate when application is made protectively before disease occurs. Use the higher rate when the disease is first observed and when the minimum number of applications are applied.
Oats	Stem Rust (Puccinia graminis f. sp avenae)	All states	230 mL	-	Harvest 4 weeks	Apply at the first sign of disease and before there is an average of over 2 pustules per tiller. Ensure thorough coverage of stems and leaves.
	Crown Rust ( <i>Puccinia</i> coronata f. sp avenae)		115 to 230 mL		Grazing 7 days	Apply after flag blade leaf is fully emerged or Z39 and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5-10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Suppression of Septoria Leaf Blotch ( <i>Leptosphaeria avenaria</i> )					Apply after flag blade is fully emerged or Z39. If infection averages 10% on the flag-2 leaf. The high rate of application gives longer period of protection than the lower rates. Use higher rates on high potential crops when conditions favour severe disease development during flowering. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
Peanuts	Early Leaf Spot ( <i>Cerecospora</i> <i>arachidicola</i> ), Late Leaf Spot ( <i>Cercosporidium</i> <i>personatum</i> )	Sth Qld, NSW, WA only	185 to 275 mL	-	14 days	This use is subject to a DMI anti- resistance strategy. Spray when disease symptoms are first observed. Apply at 14 days intervals while weather conditions favour disease. Use rates towards the higher end of the range when wet conditions prevail. Use a fungicide from a different activity group (non-DMI) after 3 consecutive sprays using REGIMENT 550 alone. Apply a maximum
	Rust ( <i>Puccinia arachidis</i> )	Sth Qld only				of 5 sprays per season. The leaves of peanuts sprayed may become darker green in colour and modified in shape. These effects will not adversely affect yield at recommended rates.

Crop	Disease Controlled	State	Rate		WHP	Critical Comments
•			Per Hectare	High volume		
Peppermint, Spearmint grown for oil production only	Mint Rust ( <i>Puccinia menthae</i> )	NSW, Vic, Tas only	230 mL	-	5 weeks	Apply 2 to 5 applications at 10 to 14 days intervals during the main disease period. DO NOT use on mint grown for the fresh market.
Perennial Ryegrass	Stem Rust ( <i>Puccinia graminis</i> ), Blind Seed Disease ( <i>Gleotinia granigena</i> )	Vic only			4 weeks	Apply at ear emergence and again at anthesis.
Plums For Prune Production	Prune Rust (Tranzschelia discolour)	NSW, Vic, SA only	-	Dilute Spraying: 15 mL/100L Concentrate Spraying: Refer to the Mixing/ Application section	1 day	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying method. <b>Curative Control:</b> Apply when the disease first occurs. Further applications should be made when the disease occurs on new growth. Do NOT make more than 5 applications to any individual tree during the season. <b>Protective treatment:</b> Spray mancozeb or zineb mixed with <b>REGIMENT</b> 550 at the full recommended rates application. <b>This use is subject to a DMI anti-resistance strategy.</b>
Pineapples	Base Rot ( <i>Thielaviopsis paradoxa</i> )	Qld, NT only	-	4.5 - 9 mL /100L	-	Preplant dip: Ensure thorough coverage by totally immersing the planting material in the dip solution. Allow 50 mL of the dip solution per plant. Apply the higher rate under conditions of high disease pressure.
Poppies (Papaver somniferum)	Leaf Smut ( <i>Entyloma fuscum</i> )	TAS only	230 mL	-	4 weeks	Usage recommended by poppy contract -companies. Apply as mid season application in the full flower/petal drop period when disease is present.
Stone Fruit	Brown Rot (Blossom Blight) (blossom phase) ( <i>Monilinia laxa</i> ) Brown Rot (blossom phase) ( <i>Monilinia fructicola</i> ) Brown Rot (fruit phase)	Vic, Tas, WA only Qld, NSW, Tas, SA, WA only Qld, NSW, Vic, Tas, SA, WA only	-	Dilute spraying: 11 mL/100L Concentrate Spraying: Refer to the Mixing/ Application section	1 day	Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. This use is subject to a DMI anti-resistance strategy. Apply at early (1 to 10%) blossom and again at full bloom. A further application is made at shuck-fall. Only 2 consecutive applications of DMI fungicides can be made during this period. The last Blossom Blight spray and the first Brown Rot (fruit phase) spray should be regarded as consecutive applications. Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying
						Imethods. Apply 3 weeks and 1 week before harvest. Only 2 consecutive applications of DMI fungicides can be made during this period. For varieties with extended harvesting periods, a third spray during the picking period may be applied if conditions are favourable for disease development.
Sugar Cane	Pineapple Disease (Ceratocystis paradoxa)	QId, NSW, WA only	-	9 mL /100L	-	Ensure thorough coverage of the cut ends of sugar cane setts
Wheat	Stripe Rust (Puccinia striiformis)	Qld, NSW, Vic, SA, WA only Tas only	115 mL or 230 mL	-	Harvest 4 weeks Grazing 7 days	Spray between stem elongation (Z30) and end of flowering (Z69) when 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rate under high infection pressure or where longer residual protection is required. Spray when 10 to 20% of leaves are infected. A repeat spray
						21 to 28 days later may be required. Use higher rate under high infection pressure or when longer residual protection is required.
	Powdery Mildew ( <i>Blumeria graminis</i> )	All States	70 to 230 mL			Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rate provides longer protection.
	Stem Rust (Puccinia graminis)		230 mL			Apply at the first sign of the disease and before there are more than 2 pustules per tiller. Ensure thorough coverage of stems and leaves.
	Septoria Tritici Blotch ( <i>Mycosphaerella</i> graminicola)		115 to 230 mL			Apply once between 70% flag leaf emergence (Z38) and early flowering (Z61). Use the higher rate under high infection pressure or where longer residual protection is required.
	Leaf Rust (Puccinia recondite f.s.p Tritici; Puccinia triticina)		70 to 230 mL			Apply after flag leaf is 70% emerged (Z38) and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5 to 10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Septoria nodorum Blotch (Phaeosphaeria nodorum)					Apply after flag leaves are around 70% emerged (Z38) if infection averages 10% on the flag-2 leaf. The high rate of application gives a longer period of protection than the lower rates. Use higher rates on higher potential crops when conditions favour severe disease development during flowering. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Yellow Spot (Pyrenophora tritici- repentis)		115 to 230 mL			Apply once between 70% flag leaf emergence (238) and early flowering (261). Use higher rate under high infection pressure or where longer residual protection is required. Apply after 70% flag leaf emergence (238) and before disease levels reach 5% on flag leaf. Higher rates give longer residual protection and often better economic returns.

spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate 4. If the dilute label rate is 25 mL/100L, then the concentrate rate becomes 4 x 25, that is 100 mL/100L of concentrate spray.

2. Your chosen concentrate spray volume: for example 500L/ba 3. The concentration factor in this example is: 4 x (i.e. 2000L/D60L = 4).

Dilute spray volume as determined above: for example 2000L/ha

being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way: FAMPLE DULY Use a sprayer designed and set up for concentrate spraying (that is a sprayer which provides water volumes less than those required to reach the point of run off) and matched to the coop Concentrate Spraying

bdd the required amount of REGINENT 550 directly to the dip and mix well. Avoid excessive contamination of the dip with organic matter.

the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volumes, and the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, point of run off. The required allote sprayer from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run off. The required allote sprayer set up and operation may also need to be changed, as the crop grows. Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout

Dilute Spraying Apricots, Plums and other Stone Fruit: Apply by high volume (dilute) sprayer or by concentrate sprayer.

Use the higher rates when applying to dense crops.

Uncloning the higher when spring the properties of the provided and stems is obtained. The object of spraying is to keep the upper 2 to 3 leaves green and function in the higher object of spraying is to keep the upper 2 to 3 leaves green and functions are made when applications are made with a cross wind of not lease than 5 knots. The higher rates when applications are made with a cross wind of not lease than 5 knots. hpiication

DO NOT graze treated turf or feed grass clipping from treated area to poultry or livestock.

Pineapples - Preplant dip

Add the required amount directly to the spray tank and mix well.

tequest. To Fundational information is listed in the safety data sheet for ProForce REGINENT 550 Fungicide is available from Indigo Specialty Products Pty Ltd on request.

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed D0 NOT induce vomiting.

**DIA TZRIA** 

out immediately with water. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing. spray wear cotion overalls buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, goggles and a disposable mist face mask to cover the mouth and no mose. When main the registrate gloves goggles if product in eyes, wash if the mouth and a mashable hat, elbow length chemical resistant gloves, goggles and a disposable mist accertance and accertance and a disposable mist accertance and accertance and accertance and a disposable mist accertance and a disposable accertance and accertance accertance accertance and accertance and accertance accertance accertance accertance accertance and accertance accertan Harminu II swallowed. Will damage eyes. Will intritate the nose and throat, and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing the SAFETY DIRECTIONS

#### purn empty containers or product.

from water sources or homes. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging 500 mm below the sunface in a disposal pit packaging to an approved waste management facility is not available bury the empty packaging 500 mm below the sunface in a disposal pit packaging to an approved waste management avaite management facility is not available bury the empty packaging 500 mm below the sunface in a disposal pit packaging to an approved waste management avaite management facility is not avaitable bury the empty packaging 500 mm below the sunface in a disposal pit packaging to an approved waste management avaite avait **JA2092ID DNA 30A9072** 

RULECTION OF LIVESTOCK

#### APVMA Approval No. 68606/121766

indigo

Indigo Speciality Products ABN 15 631 459 660 6/163-173 McEvoy Street, Alexandria NSW 2015 Ph: 0402 725 887

**NOSIO** 

OPENING OR USING CHILDREN VS BEFORE П

No warranty ( other than non-excludable statutory warranties ) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

normal conditions of use.

eller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under

CONDITIONS OF SALE

REGIMENT 550 is compatible with many commonly used orchard sprays which includes Ridomil Plus some formulations of azinphos- methyl, demeton-s-methyl, diazinon, methomyl liquid,

Mixtures with more than one of the above are not recommended. propargite, parathion, dimethoate, copper oxychloride, mancozeb, zineb and chlorothalonil.

L PROPICONAZOLE HYDROCARBON

`∂0

550

**LITUENT**:

CTIVE CONS'

ш

UNGICID

LIQUI FUNGICIDE

g/L

OLVENT: 371 GROUP 3

DO NOT enter treated area until spray has dried. PRECAUTION Re-entry Period

Peanuts, Perennial Ryegrass, in certain states and for the control of Dead Spot in Couch as specified in the

d other crops , Oats,

, Sugar Cane, Whe Queensland Blue (

certain f Fruit, S It and Qr USE.

he control of control of control of control of control of the state of the section of the sectio

Dollar

s of Bananas, ( eat and other o couch, and Sp

ases

dise

fungal

the

READ

IPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. Horoughly before opening or using this product.

vith the chemical or used containers. Istnop TON OO .neit of e PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

## in the Spring, particularly in cooler regions. These effects should be allowed to grow out before treating again. DO NOT apply more than twice per year.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER **APPROPRIATE LEGISLATION** 

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION DO NOT HARVEST FOR 1 DAY AFTER APPLICATION DO NOT HARVEST FOR 1 DAY AFTER APPLICATION Peppermint, Spearmint: Barley, Oats, Poppies, Wheat: Bananas, Stone Fruit: Perennial Ryegrass: Forage and fodder of Barley, Oats, Wheat: Pineapples, Sugar Cane: DO NOT GRAZE OR CUT FOR STOCK FOUD FOR 4 WEEKS AFTER APPLICATION NOT REQUIRED WHEN USED AS DIRECTED NOT REQUIRED WHEN USED AS DIRECTED DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION WITHHOLDING PERIOD NOT REQUIRED

#### parks and sporting areas Scerotinia homeocarpa Make a second application 14-28 days later if conditions continue to NSW, Qld, SA, WA onl Bent, Queensland Blue Couch favour disease development. Use rates toward the lower end of the range as a preventive program and against light to moderate infection. Use rates towards the higher end of the range and shorter intervals grasses in bowling greens golf greens, parks and sporting as a preventive or curative treatment when conditions are highly favourable for the disease. Couch Turf in bowling greens, parks and sporting areas Apply as a soil drench and water in immediately, ensuring thorough mixing with the soil. Spray in January to March, after renovation and recovery of active growth. Make a second application one month later where infection is severe. Spring Dead Spot NSW, Qld, Vic, SA WA only 28 mL/100 m in 150 L L*eptospaeria* spp.) 2015 later where infection is severe. DO NOT renovate treated greens until active growth has recommenced in Spring. DO NOT spray in the Spring/Summer period prior to renovation. REGIMENT 550 may cause bleaching of grass after application in late Summer/Autumn and also produce a greening effect in the following Spring. Couch may be slow to recommence active growth is the Christin particular in context regimer. These offsets abuild be **NSW** Indigo Specialty Products Pty Ltd ABN 15 631 439 660 6/163-173 McEvoy Street, Alexandria N

#### DIRECTIONS FOR USE – TURF

#### семеяел имателование в семенально в семенальности в семе

#### **EUNGICIDAL RESISTANCE WARNING**

Projectes PRGIMENT 550 Europidates, for thromologicales. For fungicide size of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the DMI group of tungicide is a member of the GMOUP **3** FUNGICIDE 550 or other Group **3** FUNGICIDE 550 or other **3** FUNGICIDE 550 or other Group **3** FUNGICIDE 550 or other **3** FUNCIDE 550 or oth

DNIXING



